

Hypertension

JOURNAL OF THE AMERICAN HEART ASSOCIATION



Learn and Live SM

Is Arterial Stiffness Related to Body Height?

Yan-Ping Liu, Tom Richart, Yan Li, Wei-Wei Zhan and Jan A. Staessen
Hypertension 2010;55:e24-e25; originally published online Apr 12, 2010;

DOI: 10.1161/HYPERTENSIONAHA.110.152553

Hypertension is published by the American Heart Association, 7272 Greenville Avenue, Dallas, TX 75214

Copyright © 2010 American Heart Association. All rights reserved. Print ISSN: 0194-911X. Online ISSN: 1524-4563

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://hyper.ahajournals.org/cgi/content/full/55/6/e24>

Subscriptions: Information about subscribing to Hypertension is online at
<http://hyper.ahajournals.org/subscriptions/>

Permissions: Permissions & Rights Desk, Lippincott Williams & Wilkins, a division of Wolters Kluwer Health, 351 West Camden Street, Baltimore, MD 21202-2436. Phone: 410-528-4050. Fax: 410-528-8550. E-mail:
journalpermissions@lww.com

Reprints: Information about reprints can be found online at
<http://www.lww.com/reprints>

Letter to the Editor

Letters to the Editor will be published, if suitable, as space permits. They should not exceed 1000 words (typed double-spaced) in length and may be subject to editing or abridgment.

Is Arterial Stiffness Related to Body Height?

To the Editor:

Experts¹ consider aortic pulse wave velocity (PWV) as the supreme index of arterial stiffness, mainly because it is an independent predictor of cardiovascular complications.² However, the noninvasive assessment of PWV is critically dependent on the measurement of the travel distance of the arterial pulse wave.³ Variability between estimates of PWV largely depends on inconsistencies in the measured travel distance.⁴ Travel distance is proportional to body height.⁵ This explains the inverse association between PWV and body height. In a recent publication in *Hypertension*, Wang et al⁶ recognized the problem and adjusted the hazard ratios expressing the risk of death related to a pulse wave reflection index for body height.

The Colin VP-1000 (Omron Healthcare) is easy to operate and is used for risk stratification, in particular in Japan and China.⁷ This device measures brachial-ankle PWV (baPWV) by means of an automated wave form analyzer. In 200 subjects (51.5% women; mean age: 51.3 years; age range: 20 to 80 years) recruited in the framework of the Flemish Study on Environment Genes and Health Outcomes,⁸ we observed a strong inverse association between baPWV and body height (Table). However, the VP-1000 device computes baPWV automatically as $(L1-L2)/T$.⁹ L1 is the distance from the heart to the ankle, L2 is the distance from the heart to the arm, and T is the time delay between the feet of the brachial and tibial arterial waveforms. The VP-1000 computes L1 and L2 from the height (H) of the subjects as $L1=0.2195 \times H - 2.0734$ and $L2=0.5643 \times H - 18.381$.⁹ For this reason, we believe that the inverse association between baPWV and body height is spurious. Leading experts in the field recently proposed other formulas to extrapolate travel distance from height.⁵ Another limitation of baPWV is that this index mixes the characteristics of the elastic aorta and subclavian arteries and the muscular brachial, femoral, and tibial arteries.

In the same Flemish Study on Environment Genes and Health Outcomes participants,⁸ we also measured aortic PWV from the length of the carotid-femoral segment and the transit time of the pulse wave. The carotid-femoral segment (cfD) was the distance between the suprasternal notch and the site of the femoral measurement (nfD) minus the distance between the site of the carotid measurement and the suprasternal notch (cnD). We measured these distances using a measuring tape. Thus, for aortic PWV, travel distance was not extrapolated from height. In contrast to baPWV, aortic PWV was not significantly correlated with body height (Table). Nevertheless, cfD ($r=0.41$), nfD ($r=0.56$), and cnD ($r=0.43$) were all closely correlated ($P<0.0001$) with body height, highlighting that even the best index of arterial stiffness not only depends on the characteristics the measured arterial segment.

We also measured the central and peripheral systolic augmentation indices.⁸ We used a high-fidelity SPC-301 micromanometer (Millar Instruments, Inc) interfaced with a laptop computer running the SphygmoCor software, version 7.1 (AtCor Medical Pty Ltd). Measurement of the systolic augmentation indices does not involve travel distance. Nevertheless, in line with our previous study of a Chinese population,¹⁰ we noticed a signifi-

Table. Association of Various Indices of Arterial Stiffness With Body Height

Characteristic	All Subjects	Women	Men
No. of subjects	200	103	97
Brachial-ankle PWV, cm/s per cm	$-5.39 \pm 2.00^*$	$-16.21 \pm 4.51^\dagger$	$-14.88 \pm 2.82^\dagger$
Carotid-femoral PWV, cm/s per cm	1.10 ± 1.32	-2.70 ± 2.36	-2.10 ± 2.51
Central AI, %/cm	$-1.26 \pm 0.18^\dagger$	$-1.48 \pm 0.36^\dagger$	$-1.24 \pm 0.37^*$
Peripheral AI, %/cm	$-1.01 \pm 0.15^\dagger$	$-1.31 \pm 0.29^\dagger$	$-1.05 \pm 0.30^\dagger$

AI indicates systolic augmentation index. Values are single regression coefficients \pm SE.

* $P<0.01$ indicates significance of the regression coefficient.

$^\dagger P<0.0001$ indicates significance of the regression coefficient.

cant inverse association between systolic augmentation and body height (Table), which can be easily explained by the faster return of the reflected wave in shorter people.

In conclusion, travel distance remains a major problem in the standardization and interpretation of PWV as the favored¹ index of arterial stiffness. The inverse association between PWV and height is overestimated if travel distance is extrapolated from height. For the same height, aortic PWV reflects only arterial stiffness, but if height is dissimilar PWV reflects both arterial properties and the difference in stature. We suggest that PWV be expressed standardized to height (and heart rate) and that analyses involving PWV as the explanatory variable should be adjusted for body height (and heart rate), as done recently by Wang et al.⁶

Disclosures

None.

Yan-Ping Liu

Department of Ultrasonography
Ruijin Hospital, Shanghai Jiaotong University School of
Medicine
Shanghai, China
Studies Coordinating Centre
Division of Hypertension and Cardiovascular Rehabilitation
Department of Cardiovascular Diseases
University of Leuven
Leuven, Belgium

Tom Richart

Studies Coordinating Centre
Division of Hypertension and Cardiovascular Rehabilitation
Department of Cardiovascular Diseases
University of Leuven
Leuven, Belgium
Department of Epidemiology
Maastricht University
Maastricht, The Netherlands

Yan Li

Center for Epidemiologic Studies and Clinical Trials and
Center for Vascular Evaluation
Ruijin Hospital
Shanghai Jiaotong University School of Medicine
Shanghai, China

Wei-Wei Zhan

Department of Ultrasonography
Ruijin Hospital, Shanghai Jiaotong University School of
Medicine
Shanghai, China

Jan A. Staessen

Studies Coordinating Centre
Division of Hypertension and Cardiovascular Rehabilitation
Department of Cardiovascular Diseases
University of Leuven
Leuven, Belgium
Department of Epidemiology
Maastricht University
Maastricht, The Netherlands

References

1. Laurent S. Surrogate measures of arterial stiffness: do they have additive predictive value or are they only surrogates of a surrogate? *Hypertension*. 2006;47:325–326.
2. Hansen TW, Staessen JA, Torp-Pedersen C, Rasmussen S, Thijs L, Ibsen H, Jeppesen J. Prognostic value of aortic pulse wave velocity as index of arterial stiffness in the general population. *Circulation*. 2006;113:664–670.
3. Weber T, Ammer M, Rammer M, Adji A, O'Rourke MF, Wassertheurer S, Rosenkranz S, Eber B. Noninvasive determination of carotid-femoral pulse wave velocity depends critically on assessment of travel distance: a comparison with invasive measurement. *J Hypertens*. 2009;27:1624–1630.
4. Rajzer MW, Wojciechowska W, Kloczek M, Palka I, Brzozowska-Kiszka M, Kawecka-Jaszcz K. Comparison of aortic pulse wave velocity measured by three techniques: Complior, SphygmoCor and Arteriograph. *J Hypertens*. 2008;26:2001–2007.
5. Vermeersch SJ, Rietzschel ER, De Buyzere ML, Van Bortel LM, Gillebert TC, Verdonck PR, Laurent S, Segers P, Boutouyrie P. Distance measurements for the assessment of carotid to femoral pulse wave velocity. *J Hypertens*. 2009;27:2377–2385.
6. Wang KL, Cheng HM, Chuang SY, Li CH, Spurgeon HA, Ting CT, Najjar SS, Lakatta EG, Yin FCP, Chou P, Chen CH. Wave reflection and arterial stiffness in the prediction of 15-year all-cause and cardiovascular mortalities: a community-based study. *Hypertension*. 2010;55:799–805.
7. Li Y, Staessen JA, Lu L, Li LH, Wang GL, Wang JG. Is isolated nocturnal hypertension a novel clinical entity? Findings from a Chinese population study. *Hypertension*. 2007;50:333–339.
8. Zebekakis PE, Nawrot T, Thijs L, Balkestein EJ, van der Heijden-Spek J, Van Bortel L, Struijker-Boudier HA, Safar ME, Staessen JA. Obesity is associated with increased arterial stiffness from adolescence until old age. *J Hypertens*. 2005;23:1839–1846.
9. Hung CS, Lin JW, Hsu CN, Chen HM, Tsai RY, Chien YF, Hwang JJ. Using brachial-ankle pulse wave velocity to associate arterial stiffness with cardiovascular risks. *Nutr Metab Cardiovasc Dis*. 2009;19:241–246.
10. Li Y, Wang JG, Dolan E, Gao PJ, Guo HF, Nawrot T, Stanton AV, Zhu DL, O'Brien E, Staessen JA. Ambulatory arterial stiffness index derived from 24-hour ambulatory blood pressure monitoring. *Hypertension*. 2006;47:359–364.